FINITENESS OF REDUCTIONS OF HECKE ORBITS

MARK KISIN, YEUK HAY JOSHUA LAM, ANANTH N.SHANKAR, AND PADMAVATHI SRINIVASAN

ABSTRACT. We prove two finiteness results for reductions of Hecke orbits of abelian varieties over local fields: one in the case of supersingular reduction and one in the case of reductive monodromy. As an application, we show that only finitely many abelian varieties on a fixed isogeny leaf admit CM lifts, which in particular implies that in each fixed dimension g only finitely many supersingular abelian varieties admit CM lifts. Combining this with the Kuga-Satake construction, we also show that only finitely many supersingular K3-surfaces admit CM lifts. Our tools include p-adic Hodge theory and group theoretic techniques.

1. INTRODUCTION

Let \overline{A} be an abelian variety over $\overline{\mathbb{F}}_p$. When \overline{A} is *ordinary*, then \overline{A} admits a canonical (CM) lift, and every isogeny from \overline{A} lifts to an isogeny in characteristic zero with source any fixed lift of \overline{A} . The aim of this paper is to show that the situation is radically different for *supersingular* abelian varieties. In fact, we prove a general theorem for all Newton strata that interpolates between the ordinary case and the supersingluar case.

The first example of an abelian variety over $\overline{\mathbb{F}}_p$ without a CM lift was given by Oort in [Oor92]. Further examples of such abelian varieties, including supersingular abelian varieties, were then constructed by Conrad, Chai and Oort in [CCO14]. We prove that such examples are in fact quite abundant.

Theorem 1.1. Only finitely many supersingular abelian varieties of a given dimension admit CM lifts.

Combining this with a refined analysis of the Kuga-Satake construction, we are able to answer a question of Ito-Ito-Koshikawa [IIK18, Remark 1.3]

Theorem 1.2. If $p \ge 5$, then only finitely many supersingular K3-surfaces over $\overline{\mathbb{F}}_p$ admit CM lifts.

We remark that the theorem above makes no mention of polarizations, so does not simply follow from the Kuga-Satake map for integral models of Shimura varieties. It instead requires an analysis of a Kuga-Satake construction at the level of p-divisible groups due to Yang [Yan19].

Our results for general Newton strata use the notion of central leaves introduced by Oort in [Oor04, Theorem 5.3]. The central leaf through \overline{A} is a closed subvariety

Date: September 14, 2021.

²⁰¹⁰ Mathematics Subject Classification. Primary 11G15, 14K10, 14K22.

Key words and phrases. Hecke orbits, CM lifts.

MARK KISIN, YEUK HAY JOSHUA LAM, ANANTH N.SHANKAR, AND PADMAVATHI SRINIVASAN

inside its Newton stratum, and essentially consists of all abelian varieties whose pdivisible group is geometrically isomorphic to that of \overline{A} . In [Oor09, Section 5], Oort computes the dimensions of central leaves and shows that they are 0-dimensional in the supersingular stratum, and equal the entire Newton stratum in the ordinary case.

Theorem 1.3. Let W be a Newton stratum in the moduli space of principally polarized abelian varieties in characteristic p. The set of points of W admitting CM lifts is contained in a finite union of central leaves.

Let A be an abelian variety over a field K, with algebraic closure \bar{K} . By the *Hecke* orbit of an abelian variety A, we mean the set of isomorphism classes of abelian varieties over \bar{K} , which are isogenous to $A_{\bar{K}}$. If A is defined over a local field, and has good reduction \bar{A} , then the image of the Hecke orbit of A in the Hecke orbit of \bar{A} is called the *reduction* of the former. We prove the following result, which is in stark contrast to the ordinary case. This is the key input for proving the CM lifting theorems stated earlier.

Theorem 1.4. Let A be an abelian variety over a characteristic zero local field, and suppose A has good supersingular reduction. Then the reduction of its Hecke orbit is finite.

This theorem answers a question posed by Poonen in an unpublished preprint, and also makes progress towards understanding the p-adic distribution of Hecke orbits. The proof of Theorem 1.4 is entirely local, and we prove an analogous theorem in the setting of p-divisible groups first.

Now suppose that K is a characteristic zero local field with ring of integers \mathcal{O}_K and residue field k. For a p-divisible group \mathscr{G} over \mathcal{O}_K , we define the Hecke orbit of \mathscr{G} and its reduction in the same way as above. In particular, the reduction is a collection of isomorphism classes of p-divisible groups over an algebraic closure of k. We also establish a finiteness theorem under a semisimplicity hypothesis on the p-adic Galois representation.

Theorem 1.5 (Theorem 2.13). Let \mathscr{G} denote a *p*-divisible group over \mathcal{O}_K , such that the *p*-adic Galois representation associated to \mathscr{G} is semisimple. Then the reduction of the Hecke orbit of \mathscr{G} is finite.

In light of Theorem 1.4 and Theorem 1.5, we make the following conjecture.

Conjecture 1.6. For any *p*-divisible group \mathscr{G} over \mathcal{O}_K , the reduction of its Hecke orbit is finite.

By Theorem 1.1 and Theorem 1.5, we know that Conjecture 1.6 holds in the case of supersingular reduction (without any semisimplicity conditions on the Galois representation), and in the case that the Galois representation is semisimple (without any condition on the Newton polygon of \mathscr{G}). We remark that unlike the situation in characteristic zero the Hecke orbit of \bar{A} can contain positive dimensional families¹. This was first observed by Moret-Bailly [MB81], who constructed a complete family of supersingular abelian surfaces over $\mathbb{P}^{1}_{\mathbb{F}_{p}}$, such that all fibers are *p*-isogenous.

¹This always happens unless the *p*-rank of the *g*-dimensional abelian variety is equal to g or g-1.

Outline of the paper: In § 2, we prove our results on finiteness of reductions of Hecke orbits. This is done using a Galois-theoretic result which relies on work of Sen and Serre. We apply this in § 3 to show the results on CM lifts of p-divisible groups and abelian varieties. Here we make crucial use of Oort's results on central leaves. Finally in § 4, we prove the finiteness result for CM lifts of supersingular K3 surfaces, by comparing the deformation theory of K3's with that of GSpin p-divisible groups.

Acknowledgements

We are very grateful to Bjorn Poonen for sharing his unpublished pre-print where we first saw the question about reductions of Hecke orbits, and for useful conversations. We are extremely grateful to Arul Shankar for several valuable conversations pertaining to CM lifts of abelian varieties. We also thank Ching-Li Chai and Ziquan Yang for valuable discussions and correspondence pertaining to CM lifts and K3 surfaces.

2. Finiteness for reductions of Hecke orbits

2.1. Let K be a field equipped with a rank 1 valuation. Throughout the paper, we will denote by \mathcal{O}_K the ring of integers of K. Fix an algebraic closure \bar{K} of K. We denote by $G_K = \operatorname{Gal}(\bar{K}/K)$ the absolute Galois group of K, and by $I_K \subset G_K$ the inertia subgroup.

In this section we suppose that K is a finite extension of \mathbb{Q}_p . We write \mathbb{F}_p for the residue field of \overline{K} . We denote by $\check{\mathbb{Q}}_p$ the maximal unramified extension of \mathbb{Q}_p , and by \check{K} the compositum of K and $\check{\mathbb{Q}}_p$.

2.2. Let $\mathscr{G}/\mathcal{O}_K$ denote a *p*-divisible group. Denote by $T_p\mathscr{G}$ the *p*-adic Tate module of \mathscr{G} , and let $\rho = \rho_{\mathscr{G}} : G_K \to \operatorname{GL}(T_p\mathscr{G})$ be the Galois representation associated to \mathscr{G} . We denote by *G* (resp. *H*) the Zariski closure of $\rho(G_K)$ (resp. $\rho(I_K)$) in $\operatorname{GL}(T_p\mathscr{G}[1/p])$. Note that, since $\rho(I_K)$ is normal in $\rho(G_K)$, *H* is normal in *G*.

Lemma 2.3. We have $[G:H] < \infty$ if and only if $[\rho(G_K):\rho(I_K)] < \infty$.

Proof. If $\rho(G_K) = \bigcup_{g \in S} g\rho(I_K)$ for some finite set S, then $\bigcup_{g \in S} gH$ is a closed set containing $\rho(G_K)$ and hence equals G. This shows that [G : H] is finite if $[\rho(G_K) : \rho(I_K)]$ is.

For the other direction, suppose that $[G : H] < \infty$. By a Theorem of Sen ([Sen73, Theorem 2]) and Serre ([Ser79, Theorem 1]), $\rho(I_K)$ is open in $H(\mathbb{Q}_p)$, and hence in $G(\mathbb{Q}_p)$, as $H(\mathbb{Q}_p)$ has finite index in $G(\mathbb{Q}_p)$. In particular, this implies that $\rho(I_K)$ is open in $\rho(G_K) \subset G(\mathbb{Q}_p)$. Since $\rho(G_K)$ is compact, this implies $\rho(G_K)/\rho(I_K)$ is discrete and compact, hence finite.

Definition 2.4. We say that two *p*-divisible groups over a finite field \mathbb{F}_q are equivalent if they become isomorphic over $\overline{\mathbb{F}}_p$.

Lemma 2.5. For any $h \ge 1$, the set of equivalence classes of p-divisible groups over \mathbb{F}_q of height h is finite.

Proof. By a result of Oort [Oor04, Corollary 1.7], there is an integer n = n(h) such that any two *p*-divisible groups over $\overline{\mathbb{F}}_p$ of height h, are isomorphic if and only if their p^n -torsion subgroups are isomorphic. In particular, the equivalence class of a *p*-divisible group \mathscr{H} over \mathbb{F}_q of height h is determined by its p^n -torsion subgroup

 $\mathscr{H}[p^n]$. Since $\mathscr{H}[p^n]$ is a finite flat group scheme over \mathbb{F}_q of order p^{nh} , there are only finitely many possibilities for $\mathscr{H}[p^n]$, and the lemma follows. \Box

2.6. Let $J(\mathscr{G})$ denote the set of isomorphism classes of *p*-divisible groups over $\mathcal{O}_{\bar{K}}$ which are isogenous to $\mathscr{G} \otimes_{\mathcal{O}_{\bar{K}}} \mathcal{O}_{\bar{K}}$. We set

$$J(\mathscr{G},\overline{\mathbb{F}}_p):=\{\mathscr{G}'\otimes_{\mathcal{O}_{K'}}\bar{\mathbb{F}}_p\mid \mathscr{G}'\in J(\mathscr{G})\},$$

the set of isomorphism classes of reductions of elements of $J(\mathscr{G})$.

Analogously, we denote by I(A) the Hecke orbit of A, namely the set isomorphisms of abelian varieties over $\mathcal{O}_{\bar{K}}$ which are isogenous to $A \otimes_{\mathcal{O}_K} \mathcal{O}_{\bar{K}}$. We then define

$$I(A, \overline{\mathbb{F}}_p) := \{ A' \otimes_{\mathcal{O}_{K'}} \overline{\mathbb{F}}_p \mid A' \in I(A) \},\$$

the reduction of the Hecke orbit of A.

We now give a Galois-theoretic criterion for the finiteness of $J(\mathscr{G}, \overline{\mathbb{F}}_p)$

Proposition 2.7. If $[G:H] < \infty$, then $J(\mathscr{G}, \overline{\mathbb{F}}_p)$ is a finite set.

Proof. By Lemma 2.3, our hypothesis implies that $\rho(I_K)$ has finite index in $\rho(G_K)$. Therefore, after replacing K by a finite extension if necessary, we may assume that $\rho(I_K) = \rho(G_K)$. Let K_{ρ} the splitting field of ρ , namely the Galois extension defined by the subgroup $\text{Ker}(\rho)$. Since $\rho(I_K) = \rho(G_K)$, K_{ρ}/K is a totally ramified extension.

Note that any isogeny of *p*-divisible groups with source \mathscr{G} can be defined over K_{ρ} . Hence if \mathscr{G}' is in $J(\mathscr{G})$, then \mathscr{G}' has a representative (again denoted \mathscr{G}') which is defined over K_{ρ} . Since K_{ρ}/K is totally ramified, the reduction $\mathscr{G}' \otimes_{\mathcal{O}_{\bar{K}}} \bar{\mathbb{F}}_p$ is defined over the residue field of K, namely \mathbb{F}_q . The finiteness of $J(\mathscr{G}, \bar{\mathbb{F}}_p)$ now follows from Proposition 2.5.

2.8. We will apply Proposition 2.7 in two cases. To explain the first of these, recall that for a *p*-divisible group \mathscr{H} over \mathbb{F}_q , its Dieudonné module $\mathbb{D}(\mathscr{H})$ is a finite free $W(\mathbb{F}_q)$ -module equipped with a semi-linear Frobenius φ . If $q = p^r$, then φ^r acts linearly on $\mathbb{D}(\mathscr{H})$, and we call this action the *q*-Frobenius Frob_{*q*} on $\mathbb{D}(\mathscr{H})$. Following Rappoport-Zink [RZ96], we say that \mathscr{H} is *decent* if the action of Frob_{*q*} on $\mathbb{D}(\mathscr{H})$ is semisimple with eigenvalues which are all rational powers of *q*. That latter condition means that for each eigenvalue α , $\alpha^m = q^n$ for some integers *m*, *n*.

The semisimplicity condition is always satisfied if $\mathscr{H} = B[p^{\infty}]$ is the *p*-divisible group arising from an abelian variety *B* over \mathbb{F}_q . Examples of a decent *p*-divisible groups include those of the form $\mathscr{H} = B[p^{\infty}]$ with *B* a supersingular abelian variety *B* over \mathbb{F}_q .

Proposition 2.9. If $\mathscr{H} = \mathscr{G} \otimes_{\mathcal{O}_K} \mathbb{F}_q$ is decent then [G:H] is finite.

Proof. Let Rep_G denote the \mathbb{Q}_p -linear Tannakian category of algebraic representations of G, and $\operatorname{Isoc}_{\mathbb{Q}_p}$ the Tannakian category of isocrystals over K_0 , the maximal absolutely unramified subfield of K. As for $\mathbb{D}(\mathscr{H})$, $\operatorname{Frob}_q := \varphi^r$ acting on $\operatorname{Isoc}_{\mathbb{Q}_p}$ is linear. Using Fontaine's functor D_{cris} , the representation $\rho : G_K \to G(\mathbb{Q}_p)$, gives rise to a functor

 $D_{\operatorname{cris}} : \operatorname{Rep}_G \to \operatorname{Isoc}_{\mathbb{Q}_p} \quad w \mapsto D_{\operatorname{cris}}(w \circ \rho).$

This functor sends $V = (T_p \mathscr{G}[1/p])^*$ to $\mathbb{D}[1/p]$, where $\mathbb{D} = \mathbb{D}(\mathscr{H})$.

Now let W be a faithful representation of G/H, viewed as a representation of G, and write $w: G \to GL(W)$. Then $\rho \circ w$ is an unramified representation, so that

the eigenvalues of Frob_q acting on $D_{\operatorname{cris}}(W)$ are p-adic units. Since W is in the Tannakian category generated by V, it follows that $D_{cris}(W)$ is in the Tannakian category generated by $\mathbb{D}[1/p]$. Hence Frob_q acting on $D_{\operatorname{cris}}(W)$ is semi-simple and each of its eigenvalues have the form $\prod_{i=1}^{n} \alpha_i^{d_i}$, where $\alpha_1, \dots, \alpha_n$ are the eigenvalues of Frob_q on $D_{\operatorname{cris}}(V)$. In particular any eigenvalue α of Frob_q acting on $D_{\operatorname{cris}}(W)$ is a rational power of q. Since α is also a p-adic unit, it follows that α is a root of unity.

It follows that Frob_q acting on $D_{\operatorname{cris}}(W)$ has finite order. This implies that for some power q' of q, W, viewed as a representation of G_K/I_K , can be identified with $(D_{\operatorname{cris}}(W) \otimes W(\mathbb{F}_{q'}))^{\varphi=1}$, with G_K/I_K acting via its action on $W(\mathbb{F}_{q'})$. In particular, we see that G_K/I_K acts on W through a finite quotient. Since $\rho(G_K)$ is dense in G, it is dense in the image of G in GL(W), and hence this image is finite. As W was a faithful representation of G/H, this proves the proposition.

Corollary 2.10. If $\mathscr{H} = \mathscr{G} \otimes_{\mathcal{O}_K} \mathbb{F}_q$ is decent, then $J(\mathscr{G}, \overline{\mathbb{F}}_p)$ is a finite set. If moroever $\mathscr{G} = A[p^{\infty}]$ for an abelian scheme A over \mathcal{O}_K with supersingular reduction, then $I(\mathscr{G}, \overline{\mathbb{F}}_p)$ is a finite set.

Proof. The first statement follows immediately Proposition 2.9 and Proposition 2.7.

For the second statement, as the special fiber of A is supersingular, we may replace K by a finite extension so that the Galois action on the prime-to-p torsion of A is through scalars (with Frobenius mapping to the scalar $q^{1/2}$ where q is the size of the residue field of K). Further, we observe that $A[p^{\infty}] \otimes_{\mathcal{O}_K} \mathbb{F}_q$ is decent as A has supersingular reduction, and hence the hypothesis of Proposition 2.7 holds. Let K_{ρ} be as in the proof of Proposition 2.7. We have that every isogeny from A is defined over K_{ρ} and hence all A' isogenous to A have reductions defined over \mathbb{F}_q , the residue field of K. By Zarhin's trick [Zar77, Theorem 4.1] there are only finitely many isomorphism classes of abelian varieties over \mathbb{F}_q , and hence $I(A, \overline{\mathbb{F}}_p)$ is a finite set. \square

2.11. Our second application of Proposition 2.7 is more indirect, and proceeds by showing that even if \mathscr{G} does not satisfy the hypothesis of Proposition 2.7 one can sometimes construct an auxiliary *p*-divisible group which does.

Lemma 2.12. Suppose that the connected component of the identity in G is reductive. Then after replacing K by a finite extension, there exists a p-divisible group $\mathscr{G}'/\mathcal{O}_K$ such that

- (1) G and G' are isomorphic over O_K
 (2) If G' (resp. H') denotes the Zariski closure of the image of G_K (resp. I_K) in $\operatorname{GL}(T_p \mathscr{G}'[1/p])$, then G' = H'.

Proof. After replacing K by a finite extension, we may assume that G is reductive, and we set T = G/H. As T is abelian, it is a torus. Let Z_G denote the center of G, and G^{der} its derived subgroup. The map $Z_G \times G^{\text{der}} \to G$ is surjective with finite kernel, so we obtain a surjective map $Z_G \to T$. This implies that there is a subtorus $T_{\text{sub}} \subset Z_G$, such that the map $T_{\text{sub}} \to T$ is an isogeny.

Let $\chi: G_K \to T(\mathbb{Q}_p)$ be the map induced by ρ , and let $\sigma \in G_K$ be a lift of the q-Frobenius. For some positive integer $m, \tau = \chi(\sigma^m)$ lifts to an element of $T_{\rm sub}(\mathbb{Q}_p)$. Thus after replacing K by a finite extension, we may assume that τ lifts to an element $\tau' \in T_{\text{sub}}(\mathbb{Q}_p)$. Since $T_{\text{sub}} \to T$ is an isogeny, the subgroup generated by τ' is bounded, so there is an unramified Galois character of the form

$$\psi: G_K \to G_K / I_K \to T_{\rm sub}(\mathbb{Q}_p); \quad \sigma \mapsto \tau'$$

Now let $C \subset Z_G(\mathbb{Q}_p)$ denote the subgroup preserving $T_p \mathscr{G}$. After replacing K by a finite extension, and so τ' by a power, we may assume that $\tau' \in C$.

The action of C on $T_p \mathscr{G}$ commutes with the action of G_K . Hence by Tate's theorem, it induces a map $C \to \operatorname{Aut} \mathscr{G}$. Since \mathscr{G} is defined over \mathcal{O}_K , for any $\gamma \in G_K/I_K$, we have a canonical isomorphism $\gamma^* \mathscr{G} \simeq \mathscr{G}$ over $\mathcal{O}_{\check{K}}$. Denote by c_γ the composite of this isomorphism and the automorphism $\psi(\gamma)^{-1} \in C$ viewed as an automorphism of \mathscr{G} (here we view ψ as a character on G_K/I_K). Then c_γ defines a descent datum on $\mathscr{G}[p^n]|_{\mathcal{O}_{\check{K}}}$, for each n. By étale descent, c_γ arises from a unique p-divisible group \mathscr{G}' over \mathcal{O}_K .

By construction, the underlying \mathbb{Z}_p -modules of $T_p\mathscr{G}$ and $T_p\mathscr{G}'$ are canonically identified, and the action of G_K on \mathscr{G}' is obtained by multiplying its action on $T_p\mathscr{G}$ by $\psi(\sigma)^{-1}$. Hence we have $H = H' \subset G' \subset G$, and the map $G' \to T$ is trivial. It follows that G' = H'.

Corollary 2.13. If the p-adic Galois representation associated to \mathscr{G} is semisimple, then the set $J(\mathscr{G}, \overline{\mathbb{F}}_p)$ is finite.

Proof. By Lemma 2.12 we can find \mathscr{G}' such that H' = G'. By Proposition 2.7 $J(\mathscr{G}', \overline{\mathbb{F}}_p)$ is a finite set. On the other hand the two *p*-divisible groups \mathscr{G} and \mathscr{G}' are isomorphic over \breve{K} and therefore $J(\mathscr{G}, \overline{\mathbb{F}}_p) = J(\mathscr{G}', \overline{\mathbb{F}}_p)$.

3. Finiteness of p-divisible groups admitting a CM lift

3.1. In this section we assume that K is a finite extension of $K_0 = W(\mathbb{F}_p)[1/p]$. A *p*-divisible group \mathscr{G} of (constant) height *h*, over any base, is said to have CM

by a commutative semisimple \mathbb{Q}_p -algebra F if there is an injective homomorphism

$$F \hookrightarrow \operatorname{End}(\mathscr{G}) \otimes \mathbb{Q}_{\mathfrak{g}}$$

such that $\dim_{\mathbb{Q}_p} F = \operatorname{height}(\mathscr{G})$. We say that \mathscr{G} is CM, or has CM if \mathscr{G} has CM by some F as above.

If \mathscr{G} is a *p*-divisible group over \mathcal{O}_K , we can form its formal group $\widehat{\mathscr{G}}$. If \mathscr{G} has CM by *F*, then $\operatorname{Lie} \widehat{\mathscr{G}} \otimes_{\mathcal{O}_K} \overline{K} \simeq \bigoplus_{\sigma} V_{\sigma}$ where σ runs over \mathbb{Q}_p -algebra maps $F \to \overline{K}$, and for $a \in F$, we have $aV_{\sigma} = \sigma(a)V_{\sigma}$. For each σ , the summand V_{σ} is either trivial, or one dimensional over \overline{K} [CCO14, Lemma 3.7.1.3]. We denote by Φ the set of σ for which V_{σ} is one dimensional, and we call Φ the CM type of \mathscr{G} .

Lemma 3.2. Let \mathscr{G} be a p-divisible group over \mathcal{O}_K with CM by F. Then there exists a finite extension K'/\mathbb{Q}_p contained in K, and a p-divisible group \mathscr{G}' over $\mathcal{O}_{K'}$ with CM by F such that $\mathscr{G}' \otimes_{\mathcal{O}_{K'}} \mathcal{O}_K$ is F-linearly isomorphic to \mathscr{G} .

Proof. This is well known. Let D be the weakly admissible module $D_{\text{cris}}(T_p\mathscr{G}[1/p])$, so that D is an $F \otimes_{\mathbb{Q}_p} K_0$ -module equipped with an injective, semi-linear Frobenius and a one step filtration $\operatorname{Fil}^1 D_K \subset D_K = D \otimes_{K_0} K$ by an $F \otimes_{\mathbb{Q}_p} K_0$ -submodule.

The filtration on D_K is induced by a cocharacter $\mu \in X_*(\operatorname{Res}_{F/\mathbb{Q}_p} \mathbb{G}_m)$. Choose K'/\mathbb{Q}_p finite and contained in K, such that μ is defined over K', and let K'_0 denote the maximal unramified subfield of K'. Then there exists a free $F \otimes_{\mathbb{Q}_p} K'_0$ -module D', an $F \otimes_{\mathbb{Q}_p} K'$ -submodule $\operatorname{Fil}^1 D'_{K'} \subset D'_{K'} = D' \otimes K'$, and an F-linear isomorphism $\iota : D' \otimes_{K'_0} K_0 \simeq D$, respecting filtrations.

Now identify D' with $F \otimes_{\mathbb{Q}_p} K'_0$. Then we can identify D with $F \otimes_{\mathbb{Q}_p} K_0$ via ι , and the Frobenius on D given by $\delta\sigma$, where $\delta \in (F \otimes_{\mathbb{Q}_p} K_0)^{\times}$ and σ denotes the Frobenius on K_0 . After possibly replacing K' by a larger field, there exists $\delta' \in (F \otimes_{\mathbb{Q}_p} K'_0)^{\times}$ such that $\delta' \delta^{-1}$ is unit. We equip D' with the Frobenius $\delta'\sigma$.

As $\delta' \delta^{-1}$ is a unit there exists $c \in (F \otimes_{\mathbb{Q}_p} K_0)^{\times}$ such that $\delta' = c^{-1} \delta \sigma(c)$. Then $c \cdot \iota$ respects Frobenius, and also respects filtrations as ι does. D' along with the Frobenius $\delta' \sigma$ and filtration $\operatorname{Fil}^1 D'_{K'} \subset D'_{K'}$ is a weakly admissible module as D is. This weakly admissible module equals $D_{\operatorname{cris}}(T_p \mathscr{G}'[1/p])$ where $\mathscr{G}'/\mathcal{O}_{K'}$ is a p-divisible group, for example by [Kis06]. The isomorphism $D'_{K_0} \simeq D_{K_0}$ induces a quasi-isogeny between the Tate-modules of \mathscr{G}' and \mathscr{G} as $\operatorname{Gal}(\bar{K}/K)$ -representations, and hence (after multiplication by a power of p) an isogeny $\mathscr{G}'_{\mathcal{O}_K} \to \mathscr{G}$. After replacing K' by a finite extension, we may assume that the kernel of this isogeny is defined over $\mathcal{O}_{K'}$ and the theorem follows.

3.3. Let \mathscr{H} be a *p*-divisible group over $\overline{\mathbb{F}}_p$. We say that \mathscr{H} admits a CM lift, if there exists a finite extension $K/W(\overline{\mathbb{F}}_p)[1/p]$, and a CM *p*-divisible group \mathscr{G} over \mathcal{O}_K , such that $\mathscr{G} \otimes_{\mathcal{O}_K} \overline{\mathbb{F}}_p$ is isomorphic to \mathscr{H} .

We remark that there is essentially no extra generality gained by considering CM lifts to more general base rings. More precisely, if R is an integral, normal, flat $W(\bar{\mathbb{F}}_p)$ -algebra, and \mathscr{G} is a CM deformation of \mathscr{H} to R, then there is a finite extension $K/W(\bar{\mathbb{F}}_p)[1/p]$, and an inclusion $\mathcal{O}_K \to R$, such that \mathscr{G} arises from a CM deformation of \mathscr{H} over \mathcal{O}_K . This can be deduced from the fact that the rigid analytic period morphism in [RZ96, §5] is étale, together with the fact that any $F \otimes_{W(\bar{\mathbb{F}}_p)} R$ -direct summand of the free, rank 1 $F \otimes_{W(\bar{\mathbb{F}}_p)} R$ -module $\mathbb{D}(\mathscr{G})(R)$ is defined over $F \otimes_{W(\bar{\mathbb{F}}_p)} \mathcal{O}_K$, for some $\mathcal{O}_K \subset R$, as above.

Theorem 3.4. Let $\mathscr{H}/\bar{\mathbb{F}}_p$ be a p-divisible group. Then, the isogeny class of \mathscr{H} contains only finitely many isomorphism classes of p-divisible groups which admit a CM lift.

Proof. Since the algebra $\operatorname{End}(\mathscr{H}) \otimes \mathbb{Q}$ has finite dimension over \mathbb{Q}_p , there are only finitely many choices for the CM algebra F. Given F, there are only finitely many choices for the CM type Φ . Thus, we may fix F and Φ , and consider only p-divisible groups in the isogeny class of \mathscr{H} , which admit a CM lift having CM by F and CM type Φ .

Let $\mathscr{G}, \mathscr{G}_1$ be such lifts, defined over some finite extension $K/W(\mathbb{F}_p)[1/p]$. By Lemma 3.2, there exists a *p*-divisible group \mathscr{G}' with CM by *F*, defined over a finite extension K'/\mathbb{Q}_p such that $\mathscr{G}' \otimes_{\mathcal{O}_{K'}} \mathcal{O}_K \simeq \mathscr{G}$. Since \mathscr{G}_1 and \mathscr{G} have the same CM type, there is an *F*-linear isogeny $\mathscr{G} \to \mathscr{G}_1$, by [CCO14, Proposition 3.7.4]. Thus $\mathscr{G}_1 \in J(\mathscr{G}')$.

Now the Zariski closure of the image of $G_{K'}$ acting on $T_p \mathscr{G}'$ is a closed subgroup of the torus $\operatorname{Res}_{F/\mathbb{Q}_p} \mathbb{G}_m$, hence is reductive. Hence $J(\mathscr{G}', \overline{\mathbb{F}}_p)$ is finite by Theorem 2.13, and the theorem follows.

3.5. Let \mathcal{A}_g denote the moduli space of principally polarized abelian varieties of dimension g. For a given Newton polygon ν we denote by $W_{\nu} \subset \mathcal{A}_{g,\bar{\mathbb{F}}_p}$ the corresponding Newton stratum. It is a locally closed subscheme.

For any $x \in \mathcal{A}_g(\bar{\mathbb{F}}_p)$, the associated *p*-divisible group \mathscr{G}_x carries a principal polarization, an isomorphism, ψ_x , of \mathscr{G}_x and its Cartier dual. The polarized central leaf through a point $x \in \mathcal{A}_g$ is the locus of points where the associated polarized *p*-divisible group is geometrically *isomorphic* to the polarized *p*-divisible group parameterized by x. Oort [Oor04, Theorem 5.3] has shown that this locus is a closed subvariety of the Newton stratum through x.

Theorem 3.6. Let S be the set of points in W_{ν} which admit CM lifts. Then S is contained in a finite number of central leaves.

Proof. Let *n* denote some large enough integer, such that $\mathcal{A}_g[n]$, the moduli space of principally polarized *g*-dimensional abelian varieties with full symplectic level *n* structure, is a fine moduli space. The notion of a polarized central leaf in $\mathcal{A}_g[n]$ through any point is defined exactly as in the case of \mathcal{A}_g , and without any reference to level structure. Then, the result for \mathcal{A}_g follows directly from the result for $\mathcal{A}_g[n]$.

Let $x \in \mathcal{W}_{\nu}(\overline{\mathbb{F}}_p) \subset \mathcal{A}_g[n](\overline{\mathbb{F}}_p)$, and let \mathscr{H}_x and \mathcal{A}_x be the associated principally polarized *p*-divisible group and abelian scheme, respectively. By [Oor04, Theorem 2.2], the set of points $y \in \mathcal{A}_g[n]$ with \mathscr{H}_y isomorphic to \mathscr{H}_x (as unpolarized *p*divisible groups) is a closed subvariety of \mathcal{W}_{ν} . By [Oor04, Theorem 3.3], this closed subvariety is a union of finitely many central leaves² in \mathcal{W}_{ν} . The result now follows immediately from Theorem 3.4.

Corollary 3.7. There are only finitely many supersingular principally polarized abelian varieties of dimension g, which admit a CM lift.

Proof. This follows from Theorem 3.6, and the fact that central leaves in the supersingular stratum are zero-dimensional ([Oor09, Section 5]). \Box

4. CM LIFTS OF SUPERSINGULAR K3-SURFACES

In this section we deduce Theorem 1.2 for CM lifts of K3-surfaces when $p \ge 5$. The weaker statement for polarized K3-surfaces is an immediate consequence of Theorem 3.6 and the theory of integral models of Shimura varieties [MP15]. Here we prove the stronger result for K3-surfaces with no reference to polarizations. The main inputs will be the Kuga-Satake construction for K3-crystals (without any reference to polarizations) due to Yang, and the crystalline Torelli theorem, due to Ogus.

4.1. Let R be a p-adically complete and separated, p-torsion free ring. Suppose that R is equipped with a lift of Frobenius σ . For $i \ge 0$, we denote by R(i), the R-module R, equipped with a Frobenius $\varphi = p^i \cdot \sigma$, and a filtration given by where $\operatorname{Fil}^2 R(2) = R(2)$, and $\operatorname{Fil}^3 R(2) = 0$.

A K3-crystal over R ([cf. [Ogu79, Definition 3.1]) is a free R-module \mathbb{L} of rank 22, endowed with a Frobenius-linear endomormophism $\varphi : \mathbb{L} \to \mathbb{L}$ and a φ -compatible, perfect, symmetric bilinear form $\langle \cdot, \cdot \rangle : \mathbb{L} \times \mathbb{L} \to R(2)$ satisfying:

(a) $p^2 \mathbb{L} \subset \varphi(\mathbb{L}).$

(b) The image of $\varphi \otimes_R R/pR$ is projective of rank 1.

When R = W(k) for a perfect field k, we also call this a K3-crystal over k.

A filtered K3-crystal over R is a K3-crystal \mathbb{L} over R, equipped with a filtration $0 = \operatorname{Fil}^3 \subset \operatorname{Fil}^2 \subset \operatorname{Fil}^1 \subset \operatorname{Fil}^0 = \mathbb{L}$ such that $\operatorname{gr}^{\bullet}\mathbb{L}$ is a projective R-module and the following conditions hold

(c) $\langle \cdot, \cdot \rangle : \mathbb{L} \otimes \mathbb{L} \to R(2)$ is strictly compatible with filtrations.

²Both of Oort's results are stated in the setting of families of *p*-divisible groups, which is the only reason we need to work with $\mathcal{A}_{g}[n]$ instead of \mathcal{A}_{g} .

(d) $\operatorname{Fil}^1 \mathbb{L} \otimes_R R/pR$ is the kernel of φ on $\mathbb{L} \otimes_R R/R$.

Note that these conditions imply that $\operatorname{Fil}^1 = \operatorname{Fil}^{2^{\perp}}$, that $\varphi(\operatorname{Fil}^2 \mathbb{L}) \subset p^2 \mathbb{L}$, and that for $i = 0, 1, 2, \operatorname{gr}^i \mathbb{L}$ has rank 1, 20, 1 respectively.

4.2. Now suppose that R = W(k) with k a perfect field, which will be either \mathbb{F}_p or a finite field in application.

A K3-crystal \mathbb{L} over k is said to be supersingular, if the slopes of φ are all 1. If \mathbb{L} is supersingular, and k is algebraically closed, then $\mathbb{L}^{\varphi=p}$ is a free \mathbb{Z}_p -module of rank 22 ([Ogu79, Theorem 3.3]), which also admits a bilinear form (which will no longer be perfect).

If $k = \mathbb{F}_q$ is a finite field, then we say a K3-crystal \mathbb{L} over k is decent if the q-Frobenius on \mathbb{L} has eigenvalues which are rational powers of q. We note that this implies the \mathbb{Z}_p -module $\mathbb{L}^{\varphi=p} \subset \mathbb{L}$ has rank 22. Note that every K3-crystal over $\overline{\mathbb{F}}_p$ admits a decent model over \mathbb{F}_q for some q, see [Kot85, §4.3]

Lemma 4.3. Let \mathbb{L} be a filtered K3-crystal over R, as above. Then the filtration on \mathbb{L} is induced by a cocharacter $\mu : \mathbb{G}_m \to \mathrm{GO}(\mathbb{L}, \langle \cdot, \cdot \rangle)$. In particular, the subgroup $P \subset \mathrm{GO}(\mathbb{L}, \langle \cdot, \cdot \rangle)$ preserving the filtration is parabolic.

Proof. Let $\mathbb{L}^2 = \operatorname{Fil}^2 \mathbb{L}$, and choose a submodule $\mathbb{L}^1 \subset \operatorname{Fil}^1$ so that $\operatorname{Fil}^1 \mathbb{L} = \mathbb{L}^2 \oplus \mathbb{L}^1$. Then $(\mathbb{L}^1)^{\perp}$ is free of rank 2 and surjects on $\operatorname{gr}^0 \mathbb{L}$, as $\langle \cdot, \cdot \rangle$ is perfect and strict for filtrations. Thus, we can choose a rank 1 direct summand $\mathbb{L}^0 \subset (\mathbb{L}^1)^{\perp}$ which maps isomorphically to $\operatorname{gr}^0 \mathbb{L}$. Then $\langle \cdot, \cdot \rangle$ induces a perfect pairing between the rank 1 subspaces $\mathbb{L}^0, \mathbb{L}^2 \subset (\mathbb{L}^1)^{\perp}$. Thus, modifying $\mathbb{L}^0 \subset (\mathbb{L}^1)^{\perp}$ we may assume that \mathbb{L}^0 is isotropic for $\langle \cdot, \cdot \rangle$.

Now $\mathbb{L} = \mathbb{L}^2 \oplus \mathbb{L}^1 \oplus \mathbb{L}^0$, and we define μ by requiring that $\mu(z)$ by z^i on \mathbb{L}^i . Since \mathbb{L}^i and \mathbb{L}^j are orthogonal for $i + j \neq 2$, $\mu(z)$ acts on $\langle \cdot, \cdot \rangle$ by z^2 . In particular μ factors through $\operatorname{GO}(\mathbb{L}, \langle \cdot, \cdot \rangle)$.

4.4. Now let k be a perfect field of characteristic p, which is either algebraically closed or an algebraic extension of \mathbb{F}_p . Let X/k denote a K3-surface. It is well known that the deformation functor of X is smooth and pro-representable and formally smooth of dimension 20 over W = W(k). Let $\operatorname{Spf} \hat{R}_X$ denote the universal deformation space of X, and let $\mathbb{X}^u \xrightarrow{\pi} \operatorname{Spf} \hat{R}$ denote the universal deformation of X. Choose a set $\{p, x_1 \dots x_{20}\}$ of elements that generate the maximal ideal of \hat{R}_X ; define σ to be the lift of the Frobenius endomorphism of $\hat{R}_X \mod p$ such that σ is the usual Frobenius on $W(\mathbb{F}_q)$, and $\sigma(x_i) = (x_i)^p$.

Then $\mathbb{L}_u = R^2 f_* \pi$ is a filtered K3-crystal over \hat{R}_X . To give a more explicit description of this filtered K3-crystal, we need the following

Lemma 4.5. Let \mathbb{L} be the K3-crystal attached to X. There exists a free \mathbb{Z}_p -module with quadratic form $(T, \langle \cdot, \cdot \rangle')$ and an isomorphism $\iota : (T, \langle \cdot, \cdot \rangle') \otimes_{\mathbb{Z}_p} W \to (\mathbb{L}, \langle \cdot, \cdot \rangle).$

Proof. The bilinear form on \mathbb{L} is self dual. It follows from the theory of nondegenerate bilinear forms over finite fields that after replacing k by at most a quadratic extension, $(\mathbb{L}, \langle \cdot, \cdot \rangle)$ has determinant a square and also admits a rank-11 isotropic subspace. Indeed, there is a unique quadratic form on \mathbb{L} (up to isomorphism) that satisfies these conditions, whence it follows that $(T, \langle \cdot, \cdot \rangle')_{\mathbb{Z}_p} \otimes W$ is isomorphic to $(\mathbb{L}, \langle \cdot, \cdot \rangle)$ where T is a rank-22 free \mathbb{Z}_p -module, and $\langle \cdot, \cdot \rangle'$ is the unique self-dual quadratic form on T that has square determinant and admits a rank-11 isotropic subspace. This weaker statement (namely, after replacing k with a quadratic extension) will actually be enough for our applications, so we only sketch the proof of the full statement of the lemma.

Choose a lift \tilde{X} of X to a smooth formal scheme over W. We will apply the theory of prismatic cohomology [BS19] to \tilde{X} , using the map $\mathfrak{S} = W[\![u]\!] \xrightarrow{u \to 0} W$, where \mathfrak{S} is equipped with the Frobenius which sends u to u^p ; this is an example of a prism. We obtain a finite free \mathfrak{S} -module \mathfrak{M} , equipped with a semi-linear Frobenius, and a bilinear pairing $\langle \cdot, \cdot \rangle_{\mathfrak{M}}$ such that

- (1) $(\mathfrak{M}, \langle \cdot, \cdot \rangle_{\mathfrak{M}}) \otimes_{\mathfrak{S}} W \simeq (\mathbb{L}, \langle \cdot, \cdot \rangle).$
- (2) There is a faithfully flat $\mathfrak{S}_{(p)}$ algebra A and an isomorphism

$$(\mathfrak{M}, \langle \cdot, \cdot \rangle_{\mathfrak{M}}) \otimes_{\mathfrak{S}} A \simeq (H^2(X_{\overline{K}}, \mathbb{Z}_p), \langle \cdot, \cdot \rangle) \otimes_{\mathbb{Z}_p} A$$

where \bar{K} is an algebraic closure of W[1/p], and $H^2(\tilde{X}_{\bar{K}}, \mathbb{Z}_p)$ is equipped with a perfect symmetric bilinear form, using Poincare duality.

Then by the Key Lemma of [Kis10, Prop. 1.3.4], one finds that there exists an isomorphism

$$(\mathbb{L}, \langle \cdot, \cdot \rangle) \simeq (H^2(X_{\bar{K}}, \mathbb{Z}_p), \langle \cdot, \cdot \rangle) \otimes W.$$

4.6. Let $\mathfrak{n} \in \operatorname{Spec} \hat{R}_X$ be the kernel of $\hat{R}_X \to W$ sending the x_i to 0, and let $\mathbb{L}_0 = \mathbb{L}_u \otimes_{\hat{R}_X} W$ be the corresponding filtered K3-crystal over W. The underlying K3-crystal of \mathbb{L}_0 is canonically identified with \mathbb{L} .

Fix an isomorphism as in Lemma 4.5. This allows us to regard $\text{GO} = \text{GO}(\mathbb{L}, \langle \cdot, \cdot \rangle)$, as a group over \mathbb{Z}_p . Choose an isomorphism $j : (\mathbb{L}_u, \langle \cdot, \cdot \rangle) \simeq (\mathbb{L}, \langle \cdot, \cdot \rangle) \otimes \hat{R}_X$ inducing the identity mod \mathfrak{n} . This gives rise to an isomorphism $\text{GO}(\mathbb{L}_u, \langle \cdot, \cdot \rangle) \simeq$ $\text{GO}(\mathbb{L}, \langle \cdot, \cdot \rangle) \otimes \hat{R}_X$, and the Frobenius maps on \mathbb{L}_u and \mathbb{L}_0 then are given by $b_u \sigma$ and $b\sigma$ for elements $b_u \in \text{GO}(\hat{R}_X[1/p])$ and $b \in \text{GO}(W[1/p])$ such that b_u specializes to b.

Proposition 4.7. The isomorphism $(\mathbb{L}_u, \langle \cdot, \cdot \rangle) \simeq (\mathbb{L}, \langle \cdot, \cdot \rangle) \otimes \hat{R}_X$ can be chosen so that

- (1) j respects filtrations.
- (2) $b_u = u \cdot b$ where $u \in U^{\text{opp}}(\hat{R}_X)$, and U^{opp} is the opposite unipotent of the parabolic $P \subset \text{GO}(\mathbb{L}_u, \langle \cdot, \cdot \rangle)$ corresponding to the filtration on \mathbb{L}_u .

Moreover, if these conditions are satisfied, the tautological map $\operatorname{Spf} \hat{R}_X \xrightarrow{u} \hat{U}^{\operatorname{opp}}$ is an isomorphism.

Proof. To show (1), we have to show that j can be chosen so that the parabolics $P \subset \operatorname{GO}(\mathbb{L}_u, \langle \cdot, \cdot \rangle)$ and $P_0 \subset \operatorname{GO}(\mathbb{L}_0, \langle \cdot, \cdot \rangle)$ corresponding to the filtrations on \mathbb{L}_u and \mathbb{L}_0 are identified. This follows from that fact that any deformation of P_0 to a parabolic in $\operatorname{GO}(\mathbb{L}_u, \langle \cdot, \cdot \rangle)$ is conjugate to the constant deformation [GP11]. The choice of such a j is unique up to conjugation by elements of $P(\hat{R}_X)$.

Next let $\mu : \mathbb{G}_m \to \text{GO}$ be a cocharacter corresponding to P. We claim that $b_u \in \text{SO}(\hat{R}_X)\sigma(\mu)(p)$. Let

$$\mathbb{L}'_u = (b_u \sigma) \mu(p^{-1}) \mathbb{L}_u = b_u \sigma(\mu)(p^{-1}) \mathbb{L}_u.$$

The conditions on the filtration in a filtered K3-crystal imply that $\mathbb{L}'_u \subset \mathbb{L}_u$, and that $\mu(p^{-1})$ acts by p^{-2} on $\langle \cdot, \cdot \rangle$. Thus for $x, y \in \mathbb{L}_u$ we have

$$\langle (b_u \sigma) \mu(p^{-1}) x, (b_u \sigma) \mu(p^{-1}) y \rangle = p^2 \sigma(\langle \mu(p^{-1}) x, \mu(p^{-1}) y \rangle) = \sigma(\langle x, y \rangle)$$

It follows that $\langle \cdot, \cdot \rangle$ is perfect on \mathbb{L}'_u , and hence $\mathbb{L}'_u = \mathbb{L}_u$. This proves the claim.

It follows that $w = b_u b^{-1} \in \mathrm{SO}(\hat{R}_X)$. Since the map $U^{\mathrm{opp}} \to P_0 \setminus \mathrm{GO}(\mathbb{L}_0, \langle \cdot, \cdot \rangle)$ is an open immersion, and w is the identity mod \mathfrak{n} , we can write $w = \lambda \cdot u$ with $u \in U^{\mathrm{opp}}(\hat{R}_X)$ and $\lambda \in P(\hat{R}_X)$ both reducing to 1 mod \mathfrak{n} . Conjugating j by an element of $P(\hat{R}_X)$ has the effect of replacing $b_u = \lambda ub$ by its σ -conjugate by the same element. Now let $m \geq 1$, and suppose that j can be chosen so that $b_u = b_u(m) = ub$ modulo \mathfrak{n}^m , so that $\lambda = \lambda(m) \equiv 1 \mod \mathfrak{n}^m$. Then,

$$\lambda^{-1}b_u\sigma(\lambda) = ub\sigma(\lambda) = u(b\sigma(\lambda)b^{-1})b.$$

Since $\lambda \equiv 1 \mod \mathfrak{n}^n$, $\sigma(\lambda) \equiv 1 \mod \mathfrak{n}^{pm}$, and hence $b_u(m+1) = \lambda^{-1}b_u(m)\sigma(\lambda) \equiv ub \mod \mathfrak{n}^{pm}$. This shows that ub is the σ -conjugate of $b_u(1)$ by the convergent product $\ldots \lambda(2)\lambda(1)$, so j can be chosen with $b_u = ub$.

It remains to show that the map $u : \operatorname{Spf} \hat{R}_X \xrightarrow{u} \hat{U}^{\operatorname{opp}}$ is an isomorphism. Given that these are both smooth 20-dimensional formal schemes over W, it suffices to prove this modulo the ideal (p, \mathfrak{m}^2) . Now let $S = \hat{R}_X/(p, \mathfrak{m}^2)$, and let \mathbb{L}_S denote filtered K3-crystal over S given by $\mathbb{L}_u|_S$. After, modifying our chosen isomorphism j by u, \mathbb{L}_S may be identified with

$$(\mathbb{L}\otimes S, u^{-1}ub\sigma(u), u \cdot (\mathrm{Fil}^{\bullet}\mathbb{L})_S, \langle \cdot, \cdot \rangle) = (\mathbb{L}\otimes S, b, u \cdot (\mathrm{Fil}^{\bullet}\mathbb{L})_S, \langle \cdot, \cdot \rangle)$$

where we have used that $\sigma(u) = 1$ in S.

Work of Nygaard-Ogus [NO85, Theorem 5.2, 5.3] implies that deformations of X to S correspond bijectively to *isotropic lifts* of Fil²L mod \mathfrak{m} . These lifts are in bijection with points of $U^{\text{opp}}(S)$ which are 1 modulo \mathfrak{m} , and hence with $\hat{U}^{\text{opp}}(S)$. This implies that u induces an isomorphism on S points, and hence is an isomorphism.

4.8. Let H denote the Clifford module associated to $(T, \langle \cdot, \cdot \rangle')$, and let $\{s_{\alpha,p}\}_{\alpha} \subset H^{\otimes}$ denote tensors whose pointwise stabilizer is the group GSpin (such tensors exist by [Kis10, Proposition 1.3.2]) of Spinor similitudes associated to $(T, \langle \cdot, \cdot \rangle')$.

Definition 4.9. A GSpin-structure on a *p*-divisible group \mathscr{H}/k is the data of an isomorphism $\iota : H \otimes W(k) \to \mathbb{D}(\mathscr{H})$, such that $\iota(s_{\alpha,p}) \in \mathbb{D}(\mathscr{H})^{\otimes}$ are Frobenius-invariant. We say that ι_1 and ι_2 are isomorphic GSpin-structures if $\iota_1(s_{\alpha,p}) = \iota_2(s_{\alpha,p})$.

4.10. We now recall a Kuga-Satake construction for K3-crystals due to Yang [Yan19, Appendix A], which associates a *p*-divisible group with GSpin structure to a K3-crystal.

Denote by \mathbb{H} the Clifford module associated to $(\mathbb{L}, \langle \cdot, \cdot \rangle)$. The isomorphism ι induces a canonical isomorphism of Clifford modules $H \otimes W(k) \to \mathbb{H}$ (where H is as in Section 4.8), which we shall also denote by ι .

Let μ be as in the proof of Proposition 4.7: a GO(L)-valued co-character, defined over W(k), that induces the mod p filtration on L. Let μ_{SO} denote the co-character of SO given by $\mu_{\text{SO}}(z) = z^{-1}\mu(z)$. The natural map

$$\operatorname{GSpin}(\mathbb{H}) \to \operatorname{SO}(\mathbb{L})$$

has kernel \mathbb{G}_m , and there is a unique lift of μ_{SO} to a cocharacter $\tilde{\mu}$ of $\operatorname{GSpin}(\mathbb{H})$ such that $\tilde{\mu}$ acts with weights 0 and 1 on \mathbb{H} . Both weights then occur with equal multiplicity. We saw in the proof of Proposition 4.7 that $b \in \operatorname{GO}(W(k))\sigma(\mu)(p)$. Hence $p^{-1}b \in \operatorname{SO}(W(k))\sigma(\mu_{SO})(p)$, and we may lift $p^{-1}b$ to an element $\tilde{b} \in$ $\operatorname{GSpin}(\mathbb{H})(W(k))\sigma(\tilde{\mu})(p)$. Then $\tilde{b}\mathbb{H} \subset \mathbb{H}$, and $\tilde{b}\mathbb{H} \not\subset p\mathbb{H}$. **Proposition 4.11** ([Yan19, Lemma A.7]). There exists a p-divisible group \mathcal{H} over k whose Dieudonné module $\mathbb{D}(\mathcal{H})$ is given by \mathbb{H} , with Frobenius acting as $\tilde{b}\sigma$.

4.12. By [HP17, Section 4.2.1], if X is supersingular, then \mathscr{H} is a supersingular p-divisible group. Since $\tilde{b} \in \operatorname{GSpin}(W[1/p])$, the tensors $s_{\alpha,0} = \iota(s_{\alpha,p}) \in \mathbb{H}^{\otimes}$ are stable by Frobenius, and therefore \mathscr{H} is equipped with a canonical GSpin-structure.

We will now use description of the universal deformation space of X to prove that \mathscr{H} admits a CM lift if X does. Write $\hat{R} = \hat{R}_X$. The opposite unipotent of $\tilde{\mu}$ in GSpin is canonically isomorphic to U^{opp} , the opposite unipotent of μ_{SO} in SO. Thus, we may regard $u \in U^{\text{opp}}(\hat{R})$ as an endomorphism of $\mathbb{H} \otimes \hat{R}$.

Let $\tilde{Fil} \subset \mathbb{H}$ denote the filtration of \mathbb{H} induced by $\tilde{\mu}$. By [Kis10, Section 1.5], the data $(\mathbb{H} \otimes \hat{R}, \tilde{Fil} \otimes \hat{R}, u \cdot (\tilde{b}\sigma))$ arises from the Dieudonné module of a *p*-divisible group $\mathscr{H}_{\hat{R}}$ over Spf \hat{R} , which deforms \mathscr{H} . Note that the tensors $s_{\alpha,0} \in \mathbb{D}(\mathscr{H}_{\hat{R}})^{\otimes}$ are Frobenius invariant and in \tilde{Fil}^0 .

Lemma 4.13. Let K/W(k)[1/p] be a finite extension, $y : \hat{R} \to \mathcal{O}_K$, a map of W(k)-algebras, and X_y (resp. \mathscr{H}_y) the K3 surface (resp. p-divisible group) over \mathcal{O}_K corresponding to y. Let $D(X_y)$ (resp. $D(\mathscr{H}_y)$) denote the weakly admissible module over K associated to X_y (resp. \mathscr{H}_y). Then there is a canonical inclusion

$$D(X_y)(1) \subset D(\mathscr{H}_y)^{\otimes}$$

compatible with filtrations and Frobenius.

Proof. The weakly admissible module $D(X_y)$ is constructed using the crystalline and de Rham cohomology of X_y , and the isomorphism between them as in [BO83, §2], and similarly for $D(\mathcal{H}_y)$. Let us briefly recall the construction.

Let \hat{R}_y denote the *PD*-completion of \hat{R} with respect to $\operatorname{Ker}(y) + p\hat{R}$. Then $\mathbb{L}_{\hat{R}_y} = \mathbb{L}_u \otimes_{\hat{R}} \hat{R}_y$ is equipped with a Frobenius and filtration. Moreover, there is a unique Frobenius equivariant map $\mathbb{L} = \mathbb{L}_0 \to \mathbb{L}_{\hat{R}_y}[1/p]$ which lifts the identity over \mathfrak{n} . It may be constructed by choosing any lift of the identity s_0 , and taking the limit $s = \lim_i \varphi^i(s_0) = \lim_i \varphi^i \circ s_0 \circ \varphi^{-i}$, which converges. This allows us to identify $\mathbb{L} \otimes_{W(k)} K$ with $\mathbb{L}_y[1/p]$, where $\mathbb{L}_y = \mathbb{L}_u \otimes_{\hat{R},y} \mathcal{O}_K$. If $K_0 \subset K$ denotes the maximal unramified subfield, this gives $D(X_y) \cong \mathbb{L} \otimes_{W(k)} K_0$ the structure of a weakly admissible module over K. Note that this identification is not in general given by the identity of \mathbb{L} . There is an analogous construction starting with $\mathbb{H}_{\hat{R}}$ in place on \mathbb{L}_u .

Now let $\mathbb{L}_u(1)$ denote the K3-crystal with underlying module \mathbb{L}_u , equipped with the Frobenius given by $p^{-1}b_u\sigma$, and the filtration given by $\operatorname{Fil}^i\mathbb{L}_u(1) = \operatorname{Fil}^{i+1}\mathbb{L}_u$. By construction, there is an inclusion $\mathbb{L}_u(1) \subset \mathbb{H}_{\hat{R}}^{\otimes}$. Applying the above construction to both sides, we obtain $D(X_y)(1) \subset D(\mathscr{H}_y)^{\otimes}$, as required. \Box

Proposition 4.14. Suppose that the K3-surface X admits a CM lift. Then so does the associated Kuga-Satake p-divisible group \mathcal{H} .

Proof. Let K, K_0 and $y : \hat{R} \to \mathcal{O}_K$, be as in the previous lemma, and denote by $G(D(X_y)(1))$ and $G(D(\mathscr{H}_y))$ the Tannakian groups of the weakly admissible modules $D(X_y)(1)$ and $D(\mathscr{H}_y)^{\otimes}$, associated to fiber functor which takes a weakly admissible module to its underlying K_0 -vector space. By construction, we have the following commutative diagram

where the vertical maps are the natural inclusions, and the top map is a surjection, which is deduced from Lemma 4.13. Since the bottom map is surjective, with kernel \mathbb{G}_m , it follows that $G(D(\mathscr{H}_y))$ is abelian if and only if $G(D(X_y)(1))$ is abelian.

Now suppose that X_y is CM. By what we just saw, it suffices to show that $G(D(X_y)(1))$, or equivalently $G(D(X_y))$, is abelian. If \bar{K} denotes an algebraic closure of K, this is equivalent to asking that the $\operatorname{Gal}(\bar{K}/K)$ action on $H^2_{\operatorname{\acute{e}t}}(X_{y,\bar{K}}, \mathbb{Q}_p)$ is abelian. This follows from work of Deligne [Del72].

Theorem 4.15. If $p \ge 5$, then only finitely many supersingular K3-surfaces over $\overline{\mathbb{F}}_p$ admit CM lifts.

Proof. Let S denote the set of supersingular K3-surfaces over $\overline{\mathbb{F}}_p$ that admit CM lifts. By Proposition 4.14, the Kuga-Satake *p*-divisible group $\mathscr{H}(X)$ associated to every $X \in S$ has a CM lift. Let S' denote the set $\{\mathscr{H}(X) : X \in S\}$. Theorem 3.4 implies that S' contains only finitely³ many $\overline{\mathbb{F}}_p$ -isomorphism classes of *p*-divisible groups, and therefore there exists a finite field \mathbb{F}_q such that every $\mathscr{H} \in S'$ admits a decent model over \mathbb{F}_q , (which we will again denote by \mathscr{H}/\mathbb{F}_q). By Lemma 4.16 below, every GSpin-structure on \mathscr{H} is defined over \mathbb{F}_q . Since the data of a *p*-divisible group \mathscr{H} along with a GSpin-structure uniquely determines the K3-crystal, it follows that the K3-crystal $\mathbb{L}(X)$ admits a decent model over \mathbb{F}_q for every $X \in S$.

By Ogus' crystalline Torelli theorem (the main result of [Ogu83]), it suffices to prove that there are only finitely many isomorphism classes of decent K3-crystals defined over any fixed finite field \mathbb{F}_q . This is a direct consequence of the discussion in [Ogu79, Section 3, Definition 3.19, Theorem 3.20] pertaining to characteristic subspaces, and therefore the result follows.

Lemma 4.16. Let \mathscr{H}/\mathbb{F}_q denote a decent p-divisible group. Then, every GSpinstructure on $\mathscr{H}_{\mathbb{F}_p}$ is defined over \mathbb{F}_q .

Proof. Let $\mathbb{H} = \mathbb{D}(\mathscr{H})$, and $\iota : H \otimes_W W(\overline{\mathbb{F}}_p) \to \mathbb{H}_{W(\overline{\mathbb{F}}_p)}$ a GSpin-structure on \mathscr{H} . Let $s_{\alpha,0} = \iota(s_{\alpha,p}) \in \mathbb{H}^{\otimes}_{W(\overline{\mathbb{F}}_p)}$. Write φ for the Frobenius on \mathbb{H} . By definition, the $s_{\alpha,0}$ are Frobenius-invariant tensors. The assumption that \mathscr{H} is decent implies that the q-power Frobenius acts on $\mathbb{H} = \mathbb{D}(\mathscr{H})$ as the scalar $q^{1/2}$. As $s_{\alpha,0}$ is $\varphi \otimes \sigma$ -invariant, it lies in the slope 0 part of $\mathbb{H}_{W(\overline{\mathbb{F}}_p)}[1/p]$. Hence, if $q = p^r$, then φ^r acts trivially on $s_{\alpha,0}$, and so σ^r acts trivially on $s_{\alpha,0}$. Hence, $s_{\alpha,0} \in \mathbb{H}^{\otimes}$.

Now consider the $W(\mathbb{F}_q)$ -scheme that represents the functor, which assigns to an $W(\mathbb{F}_q)$ -algebra R the set of isomorphisms $H \otimes R \to \mathbb{H} \otimes R$ that sends $s_{\alpha,p}$ to $s_{\alpha,0}$. This scheme is a GSpin_W -torsor as it has a point (given by the isomorphism ι) defined over $W(\overline{\mathbb{F}}_p)$. As GSpin is a connected reductive group, Lang's lemma yields that the torsor must be trivial over $W(\mathbb{F}_q)$, and hence there exists an isomorphism

³Corollary 1.2 does not follow immediately – it is a-priori possible that infinitely many nonisomorphic K3-surfaces X might yield the same Kuga-Satake p-divisible group.

 $\iota_q: H \otimes W(\mathbb{F}_q) \to \mathbb{H}$ that respects tensors. It follows that the GSpin-structure is indeed defined over \mathbb{F}_q , as claimed ⁴.

References

- [BO83] P. Berthelot and A. Ogus, F-isocrystals and de Rham cohomology. I, Invent. Math. 72 (1983), no. 2, 159–199. MR700767
- [BS19] Bhargav Bhatt and Peter Scholze, Prisms and prismatic cohomology, Preprint, available at https://arxiv.org/abs/1905.08229 (2019).
- [CCO14] Ching-Li Chai, Brian Conrad, and Frans Oort, Complex multiplication and lifting problems, Mathematical Surveys and Monographs, vol. 195, American Mathematical Society, Providence, RI, 2014. MR3137398
- [Del72] Pierre Deligne, La conjecture de Weil pour les surfaces K3, Invent. Math. 15 (1972), 206-226. MR296076
- [GP11] Philippe Gille and Patrick Polo (eds.), Schémas en groupes (SGA 3). Tome III. Structure des schémas en groupes réductifs, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], vol. 8, Société Mathématique de France, Paris, 2011. Séminaire de Géométrie Algébrique du Bois Marie 1962–64. [Algebraic Geometry Seminar of Bois Marie 1962–64], A seminar directed by M. Demazure and A. Grothendieck with the collaboration of M. Artin, J.-E. Bertin, P. Gabriel, M. Raynaud and J-P. Serre, Revised and annotated edition of the 1970 French original. MR2867622
- [HP17] Benjamin Howard and Georgios Pappas, Rapoport-zink spaces for spinor groups, Compositio Mathematica 153 (2017), no. 5, 1050–1118.
- [IIK18] Kazuhiro Ito, Tetsushi Ito, and Teruhisa Koshikawa, Cm liftings of k3 surfaces over finite fields and their applications to the tate conjecture, arXiv preprint arXiv:1809.09604 (2018).
- [Kis06] Mark Kisin, Crystalline representations and F-crystals, Algebraic geometry and number theory, 2006, pp. 459–496. MR2263197
- [Kis10] _____, Integral models for Shimura varieties of abelian type, J. Amer. Math. Soc. 23 (2010), no. 4, 967–1012. MR2669706
- [Kot85] Robert E. Kottwitz, Isocrystals with additional structure, Compositio Math. 56 (1985), no. 2, 201–220. MR809866
- [MB81] Laurent Moret-Bailly, Familles de courbes et de variétés abéliennes sur P¹. I. Descente des polarisations, 1981, pp. 109–124. Seminar on Pencils of Curves of Genus at Least Two. MR3618575
- [MP15] Keerthi Madapusi Pera, The Tate conjecture for K3 surfaces in odd characteristic, Invent. Math. 201 (2015), no. 2, 625–668. MR3370622
- [NO85] Niels Nygaard and Arthur Ogus, Tate's conjecture for K3 surfaces of finite height, Ann. of Math. (2) 122 (1985), no. 3, 461–507. MR819555
- [Ogu79] Arthur Ogus, Supersingular K3 crystals, Journées de Géométrie Algébrique de Rennes (Rennes, 1978), Vol. II, 1979, pp. 3–86. MR563467
- [Ogu83] _____, A crystalline torelli theorem for supersingular k3 surfaces, Arithmetic and geometry, 1983, pp. 361–394.
- [Oor04] Frans Oort, Foliations in moduli spaces of abelian varieties, J. Amer. Math. Soc. 17 (2004), no. 2, 267–296. MR2051612
- [Oor09] _____, Foliations in moduli spaces of abelian varieties and dimension of leaves, Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, 2009, pp. 465–501. MR2641199
- [Oor92] _____, CM-liftings of abelian varieties, J. Algebraic Geom. 1 (1992), no. 1, 131–146. MR1129842
- [RZ96] M. Rapoport and Th. Zink, Period spaces for p-divisible groups, Annals of Mathematics Studies, vol. 141, Princeton University Press, Princeton, NJ, 1996. MR1393439

⁴We note that the same proof goes through to prove that every *G*-structure on $\mathscr{H}_{\mathbb{F}_p}$ descends to \mathscr{H} whenever G/\mathbb{Z}_p is a connected reductive group. Without the connectedness assumption, the same result holds up to replacing \mathbb{F}_q with a finite extension where the degree of the extension depends only on the component set of *G*.

- [Sen73] Shankar Sen, Lie algebras of Galois groups arising from Hodge-Tate modules, Ann. of Math. (2) 97 (1973), 160–170. MR0314853
- [Ser79] Jean-Pierre Serre, Groupes algébriques associés aux modules de Hodge-Tate, Journées de Géométrie Algébrique de Rennes. (Rennes, 1978), Vol. III, 1979, pp. 155–188. MR563476
- [Yan19] Ziquan Yang, On irreducible symplectic varieties of K3^[n]-type in positive characteristic, arXiv preprint arXiv:1911.05653 (2019).
- [Zar77] Yurii Gennad'evich Zarhin, Endomorphisms of abelian varieties and points of finite order in characteristic p, Matematicheskie Zametki 21 (1977), no. 6, 737–744.

Current: M. Kisin, Department of Mathematics, Harvard University, 1 Oxford St, Cambridge, MA 02138, USA

Email address: kisin@math.harvard.edu

CURRENT: Y.H.J. LAM, DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, 1 OXFORD ST, CAMBRIDGE, MA 02138, USA

 $Email \ address: \ {\tt ylam@g.harvard.edu}$

CURRENT: A.N. SHANKAR, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, 480 LINCOLN DRIVE 213 VAN VLECK HALL, MADISON, WI 53706. Email address: ashankar@math.wisc.edu

CURRENT: P. SRINIVASAN, SCHOOL OF MATHEMATICS, UNIVERSITY OF GEORGIA, 452 BOYD GRADUATE STUDIES, 1023 D. W. BROOKS DRIVE, ATHENS, GA 30602. Email address: Padmavathi.Srinivasan@uga.edu